



Required Health Screening and Immunizations (A3.07a)

Students are required to have to have a health screening by a licensed medical provider and proof of immunizations (immunization record or titers) as recommended by the Centers for Disease Control (CDC) prior to matriculation and maintained throughout the program. The required immunizations are listed below:

Immunization	Proof of Immunization
Hepatitis B (HepB) Surface Antibody Titer (HBs)	<ul style="list-style-type: none">• Positive titer (anti-HBs at least 10mIU/mL) = immunity• Negative titer (anti-HBs less than 10mIU/mL) or never vaccinated THEN,• Give a 2-dose (Heplisav-B) or 3-dose (Engerix-B or Recombivax HB) series. Give intramuscularly (IM). For HCP who perform tasks that may involve exposure to blood or body fluids, obtain anti-HBs serologic testing 1–2 months after dose #2 (for Heplisav-B) or dose #3 (for Engerix-B or Recombivax HB).• Non-responders should be considered susceptible for Hepatitis B Virus (HBV) if their Anti-HBs remains less than 10mIU/ml after 2 vaccination series). If exposed to HepB will need Hepatitis B Immunoglobulin (HBIG).
Influenza	<ul style="list-style-type: none">• 1 dose of influenza vaccine annually.
Measles, Mumps, Rubella (MMR)-	<ul style="list-style-type: none">• Proof of vaccination or serologic evidence of MMR For healthcare personnel (HCP) born in 1957 or later without serologic evidence of immunity or prior vaccination, give 2 doses of MMR, 4 weeks apart.
Varicella	<ul style="list-style-type: none">• Proof of vaccination or serologic evidence• For HCP who have no serologic proof of immunity, prior vaccination, or diagnosis or verification of a history of varicella or herpes zoster (shingles) by a healthcare provider, give 2 doses of varicella vaccine, 4 weeks apart.



Tetanus, diphtheria, pertussis (Tdap)	<ul style="list-style-type: none">• Proof of vaccination, OR• Give 1 dose of Tdap as soon as feasible to all HCP who have not received Tdap previously, Give Td or Tdap boosters every 10 years thereafter. Give IM.
Purified Protein Derivative (PPD)	<ul style="list-style-type: none">• Students are required to have a twostep PPD for TB screening. If the screening PPD is positive, the student will be referred for follow up chest Xray and treatment if necessary. If a student has had the bacilli Calmette-Guerin (BCG) vaccine, they may receive the Tuberculin Skin Test (TST), however, they must be aware that the test may be false positive and additional tests are needed. A student may provide results of a QuantiFERON Gold or T-SPOT tests to prove immunity.
Covid-19 Vaccination	<ul style="list-style-type: none">• Pfizer-BioNTech (2 doses), or Moderna (2 doses), or Johnson & Johnson's Janssen (1 dose).